

# The AI196 antibody recognizes a SPOT-tagged recombinant protein by immunofluorescence

Wanessa Cristina Lima, Pierre Cosson

Geneva Antibody Facility, Faculty of Medicine, University of Geneva, 1 rue Michel Servet, CH-1211, Geneva, Switzerland

## Abstract

The AI196 antibody against the SPOT tag recognizes a SPOT-tagged human TAC protein by immunofluorescence in paraformaldehyde-fixed HeLa cells.

## Introduction

The SPOT tag (also called BC2 tag) is a short linear epitope derived from the human  $\beta$ -catenin protein (Uniprot #P35222), used for detection and purification of tagged proteins with the BC2-Nb antibody (Braun *et al.*, 2016). Here, we show that the AI196 recombinant antibody, derived from the BC2 nanobody, detects a SPOT-tagged human TAC protein by immunofluorescence in HeLa cells.

## Materials & Methods

**Antibodies:** The ABCD\_AI196 antibody (ABCD nomenclature, web.expasy.org/abcd/; Lima *et al.*, 2020) was produced by the Geneva Antibody Facility (www.unige.ch/medecine/antibodies/) as a mini-antibody with the antigen-binding VHH fused to a mouse IgG2A Fc. The synthesized VHH sequence (GeneArt, Invitrogen) corresponds to the sequences of the variable region of the synthetic camelid antibody BC2-Nb (Rothbauer *et al.*, 2017). HEK293 suspension cells (growing in FreeStyle™ 293 Expression Medium, Gibco #12338) were transiently transfected with the vector coding for the scFv-Fc. Supernatant (20 mg/L) was collected after 4 days.

**Antigen:** HeLa cells (growing in DMEM GlutaMAX™, Gibco #31966; supplemented with 8% Fetal Bovine Serum, Gibco #10270) cultured on glass coverslips (Menzel-Gläser, 22x22 mm) and transiently transfected 2 days before the experiment with a SPOT-tagged TAC protein (Uniprot #P01589), were used to detect the peptide tag. The SPOT epitope sequence used was PDRVRAVSHWSS and it was present in the C-terminal cytosolic domain of the fusion protein. An antibody detecting the N-terminal extracellular domain of the TAC protein (AJ519, with rabbit IgG Fc; Arsimoles *et al.*, 2020) was used as a positive control. The SPOT-tagged TAC protein is expected to be mostly present at the cell surface.

**Protocol:** The whole procedure was carried out at room temperature. Transfected HeLa cells were rinsed once with PBS, fixed with PBS + 4% paraformaldehyde (w/v) (Applichem, #A3013) for 30 min, and blocked with PBS + 40 mM ammonium chloride (NH<sub>4</sub>Cl) (Applichem, #A3661) for 5 min. Cells were then permeabilized in PBS + 0.2% saponin (w/v) (Sigma, #S7900) for 3 min, incubated in PBS + 0.2% (w/v) BSA (PBS-BSA) for 30 min, and then with the AI196 (final concentration 5 mg/L) and AJ519 antibodies (final concentration 2.5 mg/L in PBS-BSA) for 1 h. After 3 washes (10 min) with PBS-BSA, cells were incubated for 30 min in PBS-BSA with secondary goat anti-mouse IgG conjugated to AlexaFluor-647 and anti-rabbit IgG conjugated to AlexaFluor-488 (1:300, Molecular Probes, #A21235 and #A11034, respectively). After 3 washes (10 min) with PBS-BSA, cells were incubated during 10 min with DAPI (1:500, Molecular Probes, #D1306), washed twice with PBS-BSA and once with PBS, and mounted on slides (Menzel-Gläser, 76x26 mm) with Mowiol (Hoechst) + 2.5% (w/v) DABCO (Fluka, #33480). Pictures were taken using a Zeiss LSM700 confocal microscope, with a 63x Neofluar oil immersion objective.

## Results

The AI196 antibody specifically detected a signal at the plasma membrane in cells transfected with the SPOT-tagged TAC protein (Fig. 1). The signal co-localized with the signal generated by the anti-TAC AJ519 antibody (Fig. 1, arrows); the specificity of the signal was further verified by the absence of both anti-TAC and anti-SPOT stainings in the few non-transfected cells (Fig. 1, arrowheads). No staining was observed when the primary antibody was omitted (Fig. 1, No Ab).

## Conflict of interest

Pierre Cosson and Wanessa Cristina Lima are editors of the Antibody Reports journal.


## References

Arsimoles D, D'Esposito A, Gaspoz V, *et al.* The AJ519 antibody labels the human TAC/IL2RA protein by immunofluorescence. *Antibody Reports*, 2020, 3:e118. doi:10.22450/journals/abrep.2020.e118

Braun MB, Traenkle B, Koch PA, *et al.* Peptides in headlock: a novel high-affinity and versatile peptide-binding nanobody for proteomics and microscopy. *Sci Rep.* 2016; 6:19211. PMID:26791954

Lima WC, Gasteiger E, Marcatili P, Duek P, Bairoch A, Cosson P. The ABCD database: a repository for chemically defined antibodies. *Nucleic Acids Res.* 2020; 48(D1):D261-D264. PMID:31410491

Rothbauer U, Poetz O, Romer T, *et al.* Epitope tag and method for detection and/or purification of tagged polypeptides. USA; US20170137467, 2017.



**Fig. 1.** AI196 labeled the plasma membrane of HeLa cells expressing the SPOT-tagged TAC protein (in white); the signal co-localized (arrows) with the signal generated by the anti-TAC AJ519 antibody (in green); in blue, nuclei were stained with DAPI. No labelling was seen when the primary antibody was omitted, or in non-transfected cells (arrowheads). Scale bar: 20  $\mu$ m.