

# The AJ513 antibody recognizes the *Dictyostelium* p25 marker by immunofluorescence

Wanessa Cristina Lima, Pierre Cosson

Geneva Antibody Facility, Faculty of Medicine, University of Geneva, 1 rue Michel Servet, CH-1211, Geneva, Switzerland

## Abstract

The AJ513 antibody, derived from the H72 hybridoma, detects by immunofluorescence the p25-labelled compartments from *Dictyostelium discoideum*.

## Introduction

The H72 monoclonal antibody recognizes the (unidentified) p25 antigen of *D. discoideum*, used as a marker of cell surface and recycling endosomes (Ravel et al., 2001; Charette et al., 2006). Here we describe the ability of the AJ513 antibody, a single chain fragment (scFv) derived from the H72 hybridoma, to label p25 compartments by immunofluorescence.

## Materials & Methods

**Antibodies:** ABCD\_AJ513 antibody (ABCD nomenclature, web.expasy.org/abcd/) was produced by the Geneva Antibody Facility (www.unige.ch/medecine/antibodies/) as mini-antibody with the antigen-binding scFv fused to a rabbit IgG Fc. The synthesized scFv sequence (GeneArt, Invitrogen) corresponds to the sequence of the variable regions joined by a peptide linker (GGGGS)<sub>3</sub>. The sequencing of the H72 hybridoma was performed by the Geneva Antibody Facility. HEK293 suspension cells (growing in FreeStyle™ 293 Expression Medium, Gibco #12338) were transiently transfected with the vector coding for the scFv-Fc. Supernatants (~50 mg/L) were collected after 5 days.

**Antigen:** 5x10<sup>5</sup> *D. discoideum* DH1 cells, sedimented on a 22x22 mm glass coverslip (Menzel-Gläser) for 90 minutes at room temperature in HL5 medium, were used.

**Protocol:** Cells were fixed with HL5 + 4% paraformaldehyde (w/v) (Applichem, #A3013) for 30 min, and blocked with PBS + 40 mM ammonium chloride (NH<sub>4</sub>Cl) (Applichem, #A3661) for 5 min. Cells were then permeabilized in methanol at -20 °C for 2 min, washed once (5 min) with PBS, and once (15 min) with PBS + 0.2% (w/v) BSA (PBS-BSA). Cells were then incubated

for 30 min with the original mouse hybridoma H72 supernatant (dilution 1:3 in PBS-BSA) and with the reformatted scFv antibody (dilution 1:10 in PBS-BSA). After 3 washes (5, 5, 15 min) with PBS-BSA, cells were incubated for 30 min with secondary goat anti-mouse IgG conjugated to AlexaFluor-488 (hybridoma) and goat anti-rabbit IgG conjugated to AlexaFluor-647 (scFv) (1:300, Molecular Probes #A11029 and #A21245, respectively). After 3 washes (5, 5, 15 min) with PBS-BSA and one wash (5 min) with PBS, coverslips were mounted on slides (Menzel-Gläser, 76x26 mm) with Mowiol (Hoechst) + 2.5% (w/v) DABCO (Fluka, #33480). Pictures were taken using a Zeiss LSM700 confocal microscope, with a 63x Neofluar oil immersion objective.

## Results


In agreement with the original descriptions of the H72 hybridoma (Ravel et al., 2001; Charette et al., 2006), the AJ513 antibody labels the cell surface and dot-like structures in the center of the cell, characterized as recycling endosomes (Fig. 1). The staining with both antibodies appears almost indistinguishable (Fig. 1).

## References

- Charette SJ, Mercanti V, Letourneur F, Bennett N, Cosson P. A role for adaptor protein-3 complex in the organization of the endocytic pathway in *Dictyostelium*. *Traffic*. 2006;7(11):1528-38. PMID:17010123
- Ravel K, de Chassey B, Cornillon S, et al. Membrane sorting in the endocytic and phagocytic pathway of *Dictyostelium discoideum*. *Eur J Cell Biol*. 2001;80(12):754-64. PMID:11831389

## Conflict of interest

Pierre Cosson and Wanessa Cristina Lima are editors of the Antibody Reports journal.



**Fig. 1.** The H72 hybridoma and the AJ513 antibody label the cell surface and endosomal compartments in *Dictyostelium* cells. A double fluorescence staining with both antibodies was performed. No labelling was seen when the primary antibodies were omitted (No Ab). Scale bar: 10  $\mu$ m.