

Supplementary Material

Atmospheric carbon dioxide mineralisation in anthropogenically-derived carbonate deposits

John MacDonald¹, Charlotte Slaymark¹, Amanda Stubbs¹ and Marta Kalabová²

¹School of Geographical and Earth Sciences, University of Glasgow

²Department of Hydrogeology and Geothermal Resources, State Geological Institute of Dionýz Štúr

*Corresponding author: John MacDonald (john.macdonald.3@glasgow.ac.uk)

Author Note

John MacDonald

<https://orcid.org/0000-0002-8609-804X>

Charlotte Slaymark


<https://orcid.org/0000-0003-4970-6736>

Amanda Stubbs

<https://orcid.org/0000-0003-1283-2923>

Marta Kalabová

<https://orcid.org/0009-0006-9556-2293>



Supplementary Figure 1: Field and/or sample photographs for samples and/or sample locations not included in Fig. 1. a-c) other samples from the pond at Ravenscraig. d) the slag bank at Barrow, showing location of subaerial sample BW (e). f) slag-rich deposit at Clydesdale (50 cm-high rucksack for scale), showing location of subaerial sample CD2 (g). h) white-cream-orange coloured anthropogenic carbonate at Darnrigg of uncertain origin (foreground field of view ~4 m), with close up (i) showing where sample DG was obtained (2 cm-long hand lens for scale). j) field photograph showing an exposure in the flat-topped lime waste deposit at Cults (75 cm-long rucksack for scale), showing location of samples Cfl (k) and COO (l).