

The AI196 antibody recognizes a SPOT-tagged recombinant protein by immunofluorescence

Wanessa Cristina Lima, Pierre Cosson

Geneva Antibody Facility, Faculty of Medicine, University of Geneva, 1 rue Michel Servet, CH-1211, Geneva, Switzerland

Abstract

The AI196 antibody against the SPOT tag recognizes a SPOT-tagged human TAC protein by immunofluorescence in paraformaldehyde-fixed HeLa cells.

Introduction

The SPOT tag (also called BC2 tag) is a short linear epitope derived from the human β -catenin protein (Uniprot #P35222), used for detection and purification of tagged proteins with the BC2-Nb antibody (Braun *et al.*, 2016). Here, we show that the AI196 recombinant antibody, derived from the BC2 nanobody, detects a SPOT-tagged human TAC protein by immunofluorescence in HeLa cells.

Materials & Methods

Antibodies: The ABCD_AI196 antibody (ABCD nomenclature, web.expasy.org/abcd/; Lima *et al.*, 2020) was produced by the Geneva Antibody Facility (www.unige.ch/medecine/antibodies/) as a mini-antibody with the antigen-binding VHH fused to a mouse IgG2A Fc. The synthesized VHH sequence (GeneArt, Invitrogen) corresponds to the sequences of the variable region of the synthetic camelid antibody BC2-Nb (Rothbauer *et al.*, 2017). HEK293 suspension cells (growing in FreeStyle™ 293 Expression Medium, Gibco #12338) were transiently transfected with the vector coding for the scFv-Fc. Supernatant (20 mg/L) was collected after 4 days.

Antigen: HeLa cells (growing in DMEM GlutaMAX™, Gibco #31966; supplemented with 8% Fetal Bovine Serum, Gibco #10270) cultured on glass coverslips (Menzel-Gläser, 22x22 mm) and transiently transfected 2 days before the experiment with a SPOT-tagged TAC protein (Uniprot #P01589), were used to detect the peptide tag. The SPOT epitope sequence used was PDRVRAVSHWSS and it was present in the C-terminal cytosolic domain of the fusion protein. An antibody detecting the N-terminal extracellular domain of the TAC protein (AJ519, with rabbit IgG Fc; Arsimoles *et al.*, 2020) was used as a positive control. The SPOT-tagged TAC protein is expected to be mostly present at the cell surface.

Protocol: The whole procedure was carried out at room temperature. Transfected HeLa cells were rinsed once with PBS, fixed with PBS + 4% paraformaldehyde (w/v) (Applichem, #A3013) for 30 min, and blocked with PBS + 40 mM ammonium chloride (NH₄Cl) (Applichem, #A3661) for 5 min. Cells were then permeabilized in PBS + 0.2% saponin (w/v) (Sigma, #S7900) for 3 min, incubated in PBS + 0.2% (w/v) BSA (PBS-BSA) for 30 min, and then with the AI196 (final concentration 5 mg/L) and AJ519 antibodies (final concentration 2.5 mg/L in PBS-BSA) for 1 h. After 3 washes (10 min) with PBS-BSA, cells were incubated for 30 min in PBS-BSA with secondary goat anti-mouse IgG conjugated to AlexaFluor-647 and anti-rabbit IgG conjugated to AlexaFluor-488 (1:300, Molecular Probes, #A21235 and #A11034, respectively). After 3 washes (10 min) with PBS-BSA, cells were incubated during 10 min with DAPI (1:500, Molecular Probes, #D1306), washed twice with PBS-BSA and once with PBS, and mounted on slides (Menzel-Gläser, 76x26 mm) with Mowiol (Hoechst) + 2.5% (w/v) DABCO (Fluka, #33480). Pictures were taken using a Zeiss LSM700 confocal microscope, with a 63x Neofluar oil immersion objective.

Results

The AI196 antibody specifically detected a signal at the plasma membrane in cells transfected with the SPOT-tagged TAC protein (Fig. 1). The signal co-localized with the signal generated by the anti-TAC AJ519 antibody (Fig. 1, arrows); the specificity of the signal was further verified by the absence of both anti-TAC and anti-SPOT stainings in the few non-transfected cells (Fig. 1, arrowheads). No staining was observed when the primary antibody was omitted (Fig. 1, No Ab).

Conflict of interest

The authors declare no conflict of interest.

References

Arsimoles D, D'Esposito A, Gaspoz V, *et al.* The AJ519 antibody labels the human TAC/IL2RA protein by immunofluorescence. *Antibody Reports*, 2020, 3:e118. doi:10.22450/journals/abrep.2020.e118

Braun MB, Traenkle B, Koch PA, *et al.* Peptides in headlock: a novel high-affinity and versatile peptide-binding nanobody for proteomics and microscopy. *Sci Rep.* 2016; 6:19211. PMID:26791954

Lima WC, Gasteiger E, Marcatili P, Duek P, Bairoch A, Cosson P. The ABCD database: a repository for chemically defined antibodies. *Nucleic Acids Res.* 2020; 48(D1):D261-D264. PMID:31410491

Rothbauer U, Poetz O, Romer T, *et al.* Epitope tag and method for detection and/or purification of tagged polypeptides. USA; US20170137467, 2017.




Fig. 1. AI196 labeled the plasma membrane of HeLa cells expressing the SPOT-tagged TAC protein (in white); the signal co-localized (arrows) with the signal generated by the anti-TAC AJ519 antibody (in green); in blue, nuclei were stained with DAPI. No labelling was seen when the primary antibody was omitted, or in non-transfected cells (arrowheads). Scale bar: 20 μ m.